



# SEPARATION AND CHARACTERIZATION OF STEREOISOMERS OF A DINUCLEAR TRICARBONYLRHENIUM(I) COMPLEX: CHIRAL CHROMATOGRAPHY, SUPERCRITICAL VS. LIQUID

Stephen Le Garrec<sup>1</sup>, Isabelle Fabing<sup>1,2</sup>, Caroline Toppan<sup>2</sup>, Pierre Lavedan<sup>2</sup>, Mariusz Wolff<sup>3,4</sup>, Eric Benoist<sup>1</sup>, Suzanne Fery-Forgues<sup>1</sup>, Florence Bedos-Belval<sup>1</sup>

<sup>1</sup> LSPCMIB, CNRS UMR 5068, Université de Toulouse, 118 route de Narbonne, 31062, Toulouse cedex 9, France

<sup>2</sup> Institut de Chimie de Toulouse, ICT UAR 2599, Université de Toulouse, 118 route de Narbonne, 31062 Toulouse cedex 9, France


<sup>3</sup> Institut für Funktionelle Materialien and Katalyse, Währinger Straße 38-42, 1090 Wien, Austria

<sup>4</sup> Institute of Chemistry, University of Silesia in Katowice, Szkolna 9th Street, 40-006 Katowice, Poland

Chirality plays a pivotal role in modulating the biological and physicochemical properties of molecules, making it a key parameter in fields such as advanced material development, fine chemistry<sup>1</sup>, and agrochemistry.

Tricarbonylrhenium(I) complexes are gaining increasing attention due to their stability, low toxicity, and unique physicochemical properties<sup>1</sup>. Naturally chiral, these complexes exhibit enhanced structural complexity when forming dinuclear assemblies, introducing additional levels of diastereoisomerism and enantiomerism. The presence of multiple chiral centers not only enhances their potential for diverse applications<sup>2</sup> but also poses significant challenges for their separation.

In this study, we report for the first time the efficient separation<sup>3</sup> and comprehensive characterization of the two enantiomers and the meso diastereoisomer of the recently synthesized<sup>4</sup> dinuclear tricarbonylrhenium(I) complex, Bi-Re-*para*Phe. The chirality of this complex arises exclusively from its two metallic centers linked by a phenyl bridge. The performance of chiral chromatography separation is compared in both supercritical and liquid mobile phases. This methodological advancement represents a significant step forward in mastering metallic chirality, opening new perspectives for applications where precise chiral control is essential.



## References

- 1 L. C.-C. Lee, K.-K. Leung and K. K.-W. Lo, *Dalton Trans.*, **2017**, 46, 16357–16380. DOI: 10.1039/c7dt03465b
- 2 W. Yang, S. Sinha Roy, W.C. Pitts, R.L. Nelson, F.R. Fronczek, J.W. Jurss, *Inorg. Chem.*, **2018**, 57, 9564–9575. DOI: 10.1021/acs.inorgchem.8b01775.
- 3 S. Le Garrec, I. Fabing, C. Toppan, P. Lavedan, M. Wolff, E. Benoist, S. Fery-Forgues, F. Bedos-Belval, *Anal. Chim. Acta*, **2025**, 1346, 343749. DOI: 10.1016/j.aca.2025.343749.
- 4 S. Le Garrec, D. Martins-Bessa, M. Wolff, B. Delavaux-Nicot, S. Mallet-Ladeira, C. Serpentine, E. Benoist, F. Bedos-Belval, S. Fery-Forgues, *Dalton Trans.*, **2024**, 53, 16512-16529. DOI: 10.1039/D4DT01907E